



2026 AQR QCDR MEASURE SPECIFICATIONS MANUAL

2026 AQR QCDR Measure Specifications Manual

Table of Contents

Introduction	2
2025 Retired Measures	3
2026 MIPS Measures Supported	3
2026 QCDR Measures Supported	3
ePreop31: Intraoperative Hypotension (IOH) among Non-Emergent Noncardiac Surgical Cases.....	4
AQI18: Coronary Artery Bypass Graft (CABG): Prolonged Intubation – Inverse Measure	13
AQI48: Patient-Reported Experience with Anesthesia.....	15
AQI65: Avoidance of Cerebral Hyperthermia for Procedures Involving Cardiopulmonary Bypass	19
AQI71: Ambulatory Glucose Management	21
AQI79: Neuromuscular Blockade and Quantitative Train-of-Four.....	29
AQI80: Continuation of Buprenorphine for Inpatient Surgical Patients or Methadone Therapy during the Perioperative Period for Inpatient and Ambulatory Surgical Patients.....	32
AQI81: Use of Peripheral Nerve Blocks for Pain Management in Hip Fracture Patients.....	37
AQI82: Intraoperative Antibiotic Redosing in Open Cardiac Surgeries.....	39
ABG44: Low Flow Inhalational General Anesthesia	42
ABG45: Aspiration prevention in patients with gastric distension.....	45

2026 AQR QCDR Measure Specifications Manual

Introduction

Thank you for participating in the Anesthesia Quality Registry (AQR QCDR). This manual contains the specifications for all QCDR measures supported by the AQR for the 2026 reporting year. For additional details on external measures/stewards, please visit the following:

Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA) <https://aqihq.org>

ABG QCDR <https://anesthesiabg.com>

The AQR is neither responsible nor liable for changes in CMS requirements or policy. The AQR shall exercise reasonable commercial efforts to notify participants of changes in CMS requirements or policy relating to registry participation. The AQR does not represent nor warrant assurance that use of AQR's measures, data, or processes will result in participants obtaining CMS or other entity incentives or avoiding CMS or other entity penalties.

2026 AQR QCDR Measure Specifications Manual

2025 Retired Measures

Measure ID	Measure Title
424	Perioperative Temperature Management
487	Screening for Social Drivers of Health
EPREOP30	Ultrasound Guidance for Peripheral Nerve Block with Patient Experience
AQI49	Adherence to Blood Conservation Guidelines for Cardiac Operations using Cardiopulmonary Bypass (CPB) – Composite
AQI74	Continuation of Methadone Therapy during the Perioperative Period for Surgical Patients
AQI75	Continuation of Buprenorphine for Inpatient Surgical Patients
AQI76	Frailty Assessment and Intervention in Inpatient Surgical Patients
AQI77	Opioid Use Disorder Risk Screening and Referral
AQI78	Preoperative Older Adult Cognitive Assessment and Implementation of Older Adult Best Practices in Elective Surgeries
ABG42	Known or Suspected Difficult Airway Mitigation Strategies

2026 MIPS Measures Supported

Measure ID	Measure Title
130	Documentation of Current Medications in the Medical Record
145	Exposure Dose or Time for Procedures Using Fluoroscopy
155	Falls: Plan of Care
404	Anesthesiology Smoking Abstinence
430	Prevention of Post-Operative Nausea and Vomiting (PONV) - Combination Therapy
463	Prevention of Post-Operative Vomiting (POV) - Combination Therapy (Pediatrics)
477	Multimodal Pain Management

2026 QCDR Measures Supported

Measure ID	Measure Title
EPREOP31**	Intraoperative Hypotension among Non-Emergent Noncardiac Surgical Cases
AQI18	Coronary Artery Bypass Graft (CABG): Prolonged Intubation
AQI48	Patient-Reported Experience with Anesthesia
AQI65	Avoidance of Cerebral Hyperthermia for Procedures Involving Cardiopulmonary Bypass
AQI71	Ambulatory Glucose Management
AQI79	Neuromuscular Blockade and Quantitative Train-of-Four
AQI80	Continuation of Buprenorphine for Inpatient Surgical Patients or Methadone Therapy during the Perioperative Period for Inpatient and Ambulatory Surgical Patients
AQI81	Use of Peripheral Nerve Blocks for Pain Management in Hip Fracture Patients
AQI82	Intraoperative Antibiotic Redosing in Open Cardiac Surgeries – now AQI82
ABG44	Low Flow Inhalational General Anesthesia
ABG45	Aspiration prevention in patients with gastric distension

***requires EHR integration or electronically recorded MAP/blood pressure readings must be available in the required format*

2026 AQR QCDR Measure Specifications Manual

Measure Title

ePreop31: Intraoperative Hypotension (IOH) among Non-Emergent Noncardiac Surgical Cases

Measure Description: Percentage of general, neuraxial, or regional anesthesia care cases in which the mean arterial pressure (MAP) fell below 65 mmHg for a cumulative total of 15 minutes or more.

Measure Type

Intermediate Outcome

High Priority Status

Yes

Inverse Measure

Yes - A lower score indicates better quality. Note that providers are not expected to receive a score of zero on the measure, because some patients could have a MAP that falls below 65 for reasons outside a provider's control.

Instructions

This measure evaluates the proportion of cases in which the patient's MAP is below 65 mmHg for 15 minutes or more, cumulatively over the course of the surgery. The numerator condition is met when MAP is below 65 mmHg for one continuous period lasting 15 minutes or more, or if the patient has several discrete periods with a MAP below 65 mmHg that collectively sum to 15 minutes or more. Note that this measure is not intended to substitute for the clinician's judgement about managing IOH for any given patient, and for some patients the clinician may manage blood pressure using a higher or lower target MAP (e.g., a higher MAP target for patients with chronic hypertension).

To report the measure, the patient's MAP over the course of the surgery should be recorded. MAP data can be transmitted directly from the monitor to the AIMS or be entered manually into an AIMS or other electronic data capture system. If the record for a given case includes vitals pulled from the monitor and manually recorded vitals with the same time stamp, only those from the monitor will be used to score the measure.

The first blood pressure reading is defined as the anesthesia start time. The measure end time is defined as the anesthesia end time. A given blood pressure reading will be attributed to the period that runs from the time the reading was recorded to the time of either the next reading or the measure end time. If the period between a given reading and either the next reading or the measure end time lasts for longer than five minutes, the reading will only be attributed for five minutes. If the reporting clinician monitors a patient using more than one method and there are two MAPs available at the same point in time, the measure uses the invasive value for scoring the measure. The measure attributes the full case to all reporting clinicians who provide care during any portion of the case from the beginning to the end of the measurement period.

The measure excludes patients with a baseline MAP below 65 mmHg. To determine the patient's baseline MAP, the measure relies on the most recent reading taken from the preoperative holding area. If no such reading is available, the measure uses the most recent MAP taken in the operating room before induction of anesthesia.

If a clinician does not have MAP values available to report either for the baseline MAP or for measurements across the measurement period, the clinician may submit pairs of systolic and diastolic blood pressures (SBPs and DBPs) as a replacement for the MAP. The registry collecting the data will use these systolic and diastolic pressure values to calculate MAP values. Specifically, the registry will calculate MAP using the following formula: $MAP = 1/3 (SBP) + 2/3 (DBP)$ (Sesso et al. 2000).

Non-emergency surgeries include both elective and urgent surgeries.

2026 AQR QCDR Measure Specifications Manual

Because longitudinal blood pressure data can contain artifactual values (for example, inaccurate readings caused by the surgeon's leaning on the blood pressure cuff), the measure will drop MAP, SBP, and DBP readings that are likely to be artifacts. Specifically, the measure will drop individual MAP readings that meet any of the following criteria:

Documented as an artifact by the clinician

SBP \geq 300 mmHg or \leq 20 mmHg

DBP \leq 5 mmHg or DBP \geq 225 mmHg

SBP and DBP within 5 mmHg

MAP \leq 30 mmHg or \geq 250 mmHg

Measure Reporting via the Qualified Clinical Data Registry

CPT codes, patient demographics and billing data are used to identify patients who are included in the measure's denominator. Denominator eligible cases are required to be sent from an electronic reporting facility to qualify. Registry codes are used to report the numerator. Reporting clinicians who track information manually are not eligible to report the measure.

Denominator

- Unadjusted measure score: All cases in which adults (ages 18 and older) with noncardiac, non-emergency surgery requires general, neuraxial, or regional anesthesia care.
- Risk adjusted measure score: The expected number of cases in which patients have a MAP below 65 mmHg that exceeds the cumulative length of 15 minutes with noncardiac, non-emergency surgery requiring general, neuraxial, or regional anesthesia care, based on the risk adjustment model.

Denominator Criteria (Eligible Cases):

Patient aged 18 years and older

AND

Anesthesia Types: General Anesthesia, Neuraxial Anesthesia, Regional Anesthesia

AND

Patient encounter during the reporting period (CPT):

00100, 00103, 00160, 00162, 00164, 00170, 00174, 00176, 00190, 00192, 00210, 00211, 00212, 00214, 00215, 00216, 00218, 00220, 00222, 00300, 00320, 00322, 00350, 00352, 00400, 00402, 00404, 00406, 00450, 00454, 00470, 00472, 00474, 00500, 00520, 00524, 00528, 00529, 00530, 00532, 00534, 00539, 00540, 00541, 00542, 00546, 00548, 00550, 00600, 00604, 00620, 00625, 00626, 00630, 00632, 00635, 00640, 00670, 00700, 00702, 00730, 00750, 00752, 00756, 00770, 00790, 00792, 00794, 00797, 00800, 00802, 00820, 00830, 00832, 00840, 00844, 00846, 00848, 00851, 00860, 00862, 00864, 00865, 00866, 00868, 00870, 00872, 00873, 00880, 00882, 00902, 00904, 00906, 00908, 00910, 00912, 00914, 00916, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00932, 00934, 00936, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01140, 01150, 01160, 01170, 01173, 01200, 01202, 01210, 01212, 01214, 01215, 01220, 01230, 01232, 01234, 01250, 01260, 01270, 01272, 01274, 01320, 01340, 01360, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01420, 01430, 01432, 01440, 01442, 01444, 01462, 01464, 01470, 01472, 01474, 01480, 01482, 01484, 01486, 01500, 01502, 01520, 01522, 01610, 01620, 01622, 01630, 01634, 01636, 01638, 01650, 01652, 01654, 01656, 01670, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01756, 01758, 01760, 01770, 01772, 01780, 01782, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01916, 01920, 01922, 01924, 01925, 01926, 01930, 01931, 01932, 01933, 01937, 01938, 01939, 01940, 01941, 01942, 01951, 01952, 01953

2026 AQR QCDR Measure Specifications Manual

Denominator Exclusions

- **99A16** - The measure excludes patients with a baseline MAP below 65 mmHg
 - To determine the patient's baseline MAP, the measure relies on the most recent reading taken from the preoperative holding area. If no such reading is available, the measure uses the most recent MAP taken in the operating room before induction of anesthesia.
 - If a clinician does not have MAP values available to report either for the baseline MAP or for measurements across the measurement period, the clinician may submit pairs of systolic and diastolic blood pressures (SBPs and DBPs) as a replacement for the MAP. The registry collecting the data will use these systolic and diastolic pressure values to calculate MAP values. Specifically, the registry will calculate MAP using the following formula: $MAP = 1/3 (SBP) + 2/3 (DBP)$ (Sesso et al. 2000).
- **99A19** – The measure excludes patients where electronically recorded MAP/blood pressure readings are unavailable.
- **99135 CPT code** - The measure excludes surgeries where add on code 99135 (Anesthesia complicated by utilization of controlled hypotension) is listed separately in addition to the code for the primary anesthesia procedure.
- American Society of Anesthesiologists (ASA) Physical Status Classification of 1, 5 or 6
- Emergency case
- Cardiac Procedures
- Obstetric non-operative procedures
- Liver or lung transplant procedures
- Cataract procedures

Numerator

Patients who have a MAP below 65 mmHg that exceeds the cumulative length of 15 minutes.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

99A17 - MAP below 65 mmHg that exceeds the cumulative length of 15 minutes.

Performance Not Met:

99A18 - MAP does not fall below 65 mmHg for a cumulative length of 15 minutes

CBE Number: Not applicable

eCQM: Not applicable

Rationale

MAP below 60–70 mmHg among adults having non-cardiac surgery is associated with increased risk of acute kidney injury (AKI), myocardial injury, and mortality, and the risk is a function of both hypotension severity and duration (Sessler et al. 2019). Noncardiac surgery patients are at increased risk of AKI when their cumulative time below a MAP of 65 mmHg reaches or exceeds 13 minutes. When patients fall even further below this threshold (for example, MAP below 55 mmHg), even shorter durations are associated with increased risk of AKI (Salmasi et al. 2017). Among adult noncardiac surgery patients, 31.3 percent have experienced MAP below 65 mmHg for 10 minutes or longer (Bijker et al. 2007). Different approaches for managing patients' blood pressure during surgery are significantly associated with higher or lower risks of postoperative organ dysfunction, including renal dysfunction (Futier et al.2017).

Data Source: Administrative claims data; Other: Other: Claims, EHR (AIMS, patient record)

MIPS Reporting Option: Traditional MIPS, MVP

2026 AQR QCDR Measure Specifications Manual

Care Setting: Hospital; Ambulatory Surgical Center; Hospital Inpatient; Hospital Outpatient

Telehealth: No

Measure Steward: Provation Anesthesia Quality Registry (AQR QCDR)/Cleveland Clinic

Number of Multiple Performance Rates: Not applicable

Proportional Measure: No

Continuous Variable Measure: No

Ratio Measure: Yes - This is a ratio measure that will score greater than or equal to zero

Risk Adjusted: Yes

Risk adjustment: Variables incorporated into the risk adjustment model include the following:

- Age
- ASA physical status classification
- Body mass index
- Duration of surgery
- Gender

Steps for Calculating Unadjusted and Risk-Adjusted Measure Scores:

The measure is risk-adjusted to account for patient-level and case-level risk factors that affect the probability of IOH that are outside of an anesthesia provider's control. The risk adjustment model calculates the likelihood that a given case would result in IOH based on patient factors; the risk-adjusted measure then scores a clinician by comparing observed instances of IOH to the expected number of IOH cases for that clinician, given the characteristics of their patient population. Clinicians with more observed cases of IOH than expected would receive a higher (worse) score than those with fewer observed cases of IOH than expected.

Use the following steps to calculate clinician-level unadjusted and risk-adjusted measure scores. Note that this measure is specified at the individual clinician-level, but those wishing to report this measure at the group level can follow the calculation steps below but perform those calculations at the group rather than clinician-level (e.g., identifying measure denominator cases associated with the provider group).

1. First, *clean the data* to be used in calculating the measure scores. Check for missing or implausible values for key variables and drop artefactual blood pressure readings from the longitudinal blood pressure data.
2. Apply the measure logic to all cases occurring during the measurement period to identify all cases meeting *the denominator criteria*, all cases *excluded from the denominator*, and all cases meeting *the numerator criteria* (i.e., cases with IOH).
3. Calculate a *clinician-level unadjusted measure score*. This score is a percentage, with the numerator defined as all numerator cases associated with the clinician, and the denominator defined as all denominator cases (minus excluded cases) associated with the clinician.
4. Apply the risk adjustment model to calculate *the predicted probability that a given case would meet the numerator criteria* (i.e., result in IOH). The model use logistic regression to calculate the log-odds that a given case will result in IOH based on patient- and case-level factors. Apply the model to all cases that meet the denominator criteria and that are not excluded from the denominator. Transform the case-level log-odds into case-level predicted probabilities. To do so, exponentiate the log-odds to first transform it to odds. Then transform the odds to probability by taking the odds divided by 1 plus the odds.

2026 AQR QCDR Measure Specifications Manual

5. Calculate a *clinician-level expected number of IOH cases*. For a given clinician, take the sum of the predicted probabilities for all denominator cases associated with the clinician (minus exclusions). This sum represents the total number of cases for the clinician that are expected to result in IOH, given the risk level of his or her patients.
6. Calculate a *risk-adjusted score* for each clinician. The score is the ratio of the clinician's total count of cases meeting the numerator criteria to the expected number of IOH cases, among cases that meet denominator criteria for that clinician.
7. (Optional) Transform the risk-adjusted score for each clinician *into a percentage*. Note that performing this transformation is not necessary to calculate the measure, but individual sites may find that representing the scores as a percentage may be helpful for communicating with providers about their measure score. To do so, multiply each clinician's risk adjusted score from Step 6 (the observed to expected ratio) by the average unadjusted IOH measure score for the larger unit within which clinicians are being compared, for example, a group practice, hospital department, or national reporting program. This transformation may make the risk-adjusted score more easily interpretable, although it is not a true percentage generated from the ratio of numerator and denominator, and it can result in "percentages" greater than 100%.

The remainder of this document describes each of these steps in detail.

Step 1: Clean the data to be used in measure score calculation

This section described the recommended steps for cleaning the data to be included in the measure score calculation. It identifies checks to run on the data, but in most cases, it does not proscribe a specific approach for cleaning the data, leaving that determination to each individual site.

1. Check for missing values of any of the risk adjustment variables (age, gender, ASA status, BMI, surgery length); the risk adjustment model requires that all covariates are non-missing for each case. Determine how best to address missing values (e.g., impute them, or drop the case if there are few cases with missing values).
2. Check for implausible values for the risk adjustment variables. Determine how best to address them (e.g., correct them if possible, or drop the case if there are few implausible values).
3. Check for implausible values for the timestamp variables. For example, anesthesia start time and induction time should always occur before anesthesia end time. Determine how best to address implausible timelines (e.g., correct them if possible, or drop the case if there are few implausible timelines).
4. Drop artefactual blood pressure readings from longitudinal blood pressure data. See Guidelines section above for details.

Step 2 Apply measure logic to identify denominator cases, denominator exclusions, and numerator cases

This section describes the steps used to apply the measure logic to each case included in the measure's initial population. See specifications above and attached measure flow diagram for more detailed guidance on applying measure logic, including definitions of all key parameters.

1. Run the measure on all anesthesia cases during the measurement period, representing a full calendar year.
2. Apply the initial population criteria to each case (see Initial Population section above for definitions for key parameters), and remove cases from the population if any of the below scenarios applies:
 - a. Patient is under 18 years of age
 - b. Case is an emergency surgery
 - c. Case does not include general anesthesia, neuraxial, or regional anesthesia care
3. Use the cases in the initial population as *the denominator cases*.
4. *Apply denominator exclusion criteria* to the denominator cases (see Denominator Exclusions section above for definitions of key parameters), and exclude cases if any of the below scenarios applies:
 - a. Case has ASA Physical Status Classification of 1, 5 or 6
 - b. Patient has baseline MAP below 65 mmHg
 - c. Case includes induced hypotension
5. For each denominator case not excluded from the measure, apply the numerator criteria. Calculate the cumulative duration in which the patient's MAP was below 65 mmHg from anesthesia start time to anesthesia end. If this duration reaches or exceeds 15 minutes, assign the case to *the numerator population*. Otherwise, do not assign the case to the numerator population.

2026 AQR QCDR Measure Specifications Manual

Step 3: Calculate the clinician-level unadjusted measure score

This section describes the steps for calculating each clinician's unadjusted score on the IOH measure.

1. For a given clinician, identify all cases the clinician is associated with that are included in the measure denominator.
2. Calculate the clinician's unadjusted score on the measures using the following equation:

$$IOH_{Unadjusted} = \frac{(Sum\ of\ numerator\ cases)}{(Sum\ of\ denominator\ cases) - (Sum\ of\ denominator\ exclusion\ cases)}$$

Step 4: Apply risk adjustment model to calculate predicted probability of IOH

After calculating the unadjusted score, the next step is to apply the risk adjustment logistic regression model to each denominator case to determine the case's predicted probability of inclusion in the numerator population (i.e., of IOH occurring) given the case mix. The model includes five risk adjustment variables that may have an association with risk of IOH based on the clinical literature, input from experts during development of the measure, results from measure testing, or a combination of these factors. The risk adjustment variables include the patient's age, the ASA Physical Status Classification for the case, the patient's body mass index (BMI), the duration of the surgery, and the patient's gender. These variables were selected because they are associated with IOH but are outside the control of the clinician. In the model, these categorical variables with k categories are transformed into (k-1) variables with two levels.

1. Apply the risk adjustment model to each case that is part of the denominator population and that has not been excluded.
2. The risk adjustment model is a logistic regression model with the following form:

$$\begin{aligned} \text{logit}(IOH) = & \beta_0 + \beta_1 * Age + \beta_2 * ASA_2 + \beta_3 * ASA_4 + \beta_4 * BMI + \beta_5 * Surg_Length_Cat_60-119 + \beta_6 \\ & * Surg_Length_Cat_120-179 + \beta_7 * Surg_Length_Cat_180-239 + \beta_8 * Surg_Length_Cat_240-299 + \beta_9 \\ & * Surg_Length_Cat_300- + \beta_{10} * Female_1 \end{aligned}$$

Where:

β_0 = the intercept term of the logistic regression

β_1 = the coefficient for age

Age = the age in years of the patient at the time of surgery (in years)

β_2 = the coefficient for ASA physical status classification being 2.

ASA_2 = a binary variable indicating whether the ASA physical status classification of the case is 2, with *ASA_2*=1 for cases in which the ASA physical status classification is 2, and *ASA_2*=0 for cases in which it is not 2.

β_3 = the coefficient for ASA physical status classification being 4.

ASA_4 = a binary variable indicating whether the ASA physical status classification of the case is 4, with *ASA_4*=1 for cases in which the ASA physical status classification is 4, and *ASA_4*=0 for cases in which it is not 4.

β_4 = the coefficient for body mass index (BMI)

BMI = the BMI of the patient at the time of surgery

β_5 = the coefficient for the duration of surgery being between 60 and 119 minutes

Surg_Length_Cat_60-119 = a binary variable indicating whether the duration of surgery from anesthesia start time to anesthesia end time was between 60 and 119 minutes, with *Surg_Length_Cat_60-119* = 1 for surgeries that met this criteria and *Surg_Length_Cat_60-119* = 0 for surgeries that did not meet this criteria.

β_6 = the coefficient for the duration of surgery being between 120 and 179 minutes

Surg_Length_Cat_120-179 = a binary variable indicating whether the duration of surgery from anesthesia start time to anesthesia end time was between 120 and 179 minutes, with *Surg_Length_Cat_120-179* = 1 for surgeries that met this criteria and *Surg_Length_Cat_120-179* = 0 for surgeries that did not meet this criteria.

β_7 = the coefficient for the duration of surgery being between 180 and 239 minutes

Surg_Length_Cat_180-239 = a binary variable indicating whether the duration of surgery from anesthesia start time to anesthesia end time was between 180 and 239 minutes, with *Surg_Length_Cat_180-239* = 1 for surgeries that met this criteria and *Surg_Length_Cat_180-239* = 0 for surgeries that did not meet this criteria.

2026 AQR QCDR Measure Specifications Manual

- β_8 = the coefficient for the duration of surgery being between 240 and 299 minutes
 Surg_Length_Cat_240–299 = a binary variable indicating whether the duration of surgery from anesthesia start time to anesthesia end time was between 240 and 299 minutes, with Surg_Length_Cat_240–299 = 1 for surgeries that met this criteria and Surg_Length_Cat_240–299 = 0 for surgeries that did not meet this criteria.
- β_9 = the coefficient for the duration of surgery being 300 minutes or longer
 Surg_Length_Cat_300– = a binary variable indicating whether the duration of surgery from anesthesia start time to anesthesia end time was 300 minutes or longer, with Surg_Length_Cat_300– = 1 for surgeries that met this criteria and Surg_Length_Cat_300– = 0 for surgeries that did not meet this criteria.
- β_{10} = the coefficient for the gender of the patient
 Female_1 = a binary variable indicating the gender of the patient, with Female_1 = 1 for female and 0 for male.

See Table 1 for the values of the constant and the regression coefficients.

Table 1: Parameters for risk adjustment model for the intraoperative hypotension quality measure

Parameter	Value
β_0 : Constant/Intercept	-1.576
β_1 : Coefficient 1: Age	-0.008
β_2 : Coefficient 2: ASA_2	0.157
β_3 : Coefficient 3: ASA_4	0.529
β_4 : Coefficient 4: BMI	-0.018
β_5 : Coefficient 5: Surg_Length_Cat_60–119	1.316
β_6 : Coefficient 6: Surg_Length_Cat_120–179	1.734
β_7 : Coefficient 7: Surg_Length_Cat_180–239	1.936
β_8 : Coefficient 8: Surg_Length_Cat_240–299	2.235
β_9 : Coefficient 9: Surg_Length_Cat_300+	2.879
β_{10} : Coefficient 10: Female1	0.173

- The model calculates the log-odds of each case developing IOH, given the risk factors for the given patient and case. Next, transform the case-level log-odds into case-level predicted probabilities. To do so, exponentiate the log-odds to first transform it to odds. Then transform the odds to probability by taking the odds divided by 1 plus the odds. Predicted probabilities can range from 0.00 to 1.00. Values closer to 1.00 represent a higher likelihood that the case would result in IOH. The predicted probability (denoted as $IOH_{expected}$) can be presented as:

$$IOH_{expected} = \frac{e^{\text{logit}(IOH)}}{1 + e^{\text{logit}(IOH)}}$$

Where, logit(IOH) is defined in Step 4.2.

Step 5: Calculate the clinician-level expected number of IOH cases

Next, determine each clinician's expected number of IOH cases based on the risk-adjustment model by summing the case-level predicted probabilities.

- For a given clinician, identify all cases the clinician is associated with that are included in the measure denominator and that have not been excluded.
- Calculate the clinician's expected number of IOH cases by summing all of the predicted probabilities of IOH for all of the denominator cases.

Step 6: Calculate the clinician-level risk-adjusted measure score

2026 AQR QCDR Measure Specifications Manual

After computing an observed and expected number of IOH cases for each clinician, the measure uses those two values as inputs for the risk-adjusted score.

1. For a given clinician, use the observed and expected number of IOH cases to calculate the risk-adjusted score. The observed number of cases is the numerator from the equation in Step 3, and the expected number of cases is the sum calculated in Step 5. Calculate the risk-adjusted score as follows:

$$IOH_{Adjusted} = \frac{(Sum\ of\ numerator\ cases)}{(Sum\ of\ expected\ IOH\ cases)}$$

The resulting score will be a ratio. A score of 1 indicates the clinician had the number of IOH cases we would expect, based on their case mix. Scores less than 1 indicate the clinician had fewer IOH cases than predicted, meaning they are performing better than expected for their case mix. Scores greater than 1 indicate the clinician had more cases of IOH than predicted, meaning they are performing worse than expected given their case mix.

Step 7 (optional): Transform risk-adjusted measure score into a percentage

To make the risk-adjusted scores more easily interpretable, the clinician-level ratios calculated in Step 6 can be multiplied by the overall unadjusted performance rate on the measure to transform them into percentages. Note that performing this transformation is not necessary to calculate the measure, but individual sites may find that representing the scores as a percentage may be helpful for communicating with providers about their measure score. To do so, multiply each clinician's risk adjusted score from Step 6 (the observed to expected ratio) by the average unadjusted IOH measure score for the larger unit within which clinicians are being compared, for example, a group practice, hospital department, or national reporting program. This transformation may make the risk-adjusted score more easily interpretable, although it is not a true percentage generated from the ratio of numerator and denominator, and it can result in "percentages" greater than 100%.

$$IOH_{Adjusted} = \frac{(Sum\ of\ numerator\ cases)}{(Sum\ of\ expected\ IOH\ cases)} * Overall_rate$$

References

- Anesthesia Quality Institute. "2018 AQI NACOR Data Element Conceptual Definitions, Version 3.0." Schaumburg, IL: Anesthesia Quality Institute, January 2018. Available at http://www.aqihq.org/files/AQI_NACOR_DATA_ELEMENT_DEFINITIONS_v3%202018_FINAL.pdf. Accessed March 14, 2019.
- Bijker, Jilles, Wilton A. van Klei, Teus H. Kappen, Leo van Wolfswinkel, Karel G.M. Moons, and Cor J. Kalkman. "Incidence of Intraoperative Hypotension as a Function of the Chosen Definition." *Anesthesiology*, vol. 107, no. 2, 2007, pp. 213–220. doi:10.1097/01.anes.0000270724.40897.8e.
- Futier, E., J.-Y. Lefrant, P.-G. Guinot, T. Godet, E. Lorne, P. Cuvillon, S. Bertran, M. Leone, B. Pastene, V. Piriou, S. Molliex, J. Albanese, J. Julia, B. Tavernier, E. Imhoff, J. Bazin, J. Constantin, B. Pereira, and S. Jaber. "Effect of Individualized Versus Standard Blood Pressure Management Strategies on Postoperative Organ Dysfunction Among High-Risk Patients Undergoing Major Surgery: A Randomized Clinical Trial." *JAMA*, vol. 318, no. 14, 2017, pp. 1346–1357. doi:10.1001/jama.2017.14172.
- Salmasi, Vafi, Kamal Maheshwari, Dongsheng Yang, Edward J. Mascha, Asha Singh, Daniel I. Sessler, and Andrea Kurz. "Relationship Between Intraoperative Hypotension, Defined by Either Reduction from Baseline or Absolute Thresholds, and Acute Kidney and Myocardial Injury After Noncardiac Surgery: A Retrospective Cohort Analysis." *Anesthesiology*, vol. 126, no. 1, 2017, pp. 47–65. doi: 10.1097/ALN.0000000000001432.
- Sessler, Daniel I., Joshua A. Bloomstone, Solomon Aronson, Colin Berry, Tong J. Gan, John A. Kellum, James Plumb, Monty G. Mythen, Michael P. W. Grocott, Mark R. Edwards, Timothy E. Miller, and the Perioperative Quality Initiative-3 Workgroup. "Perioperative Quality Initiative Consensus Statement on Intraoperative Blood Pressure, Risk and Outcomes for Elective Surgery." *British Journal of Anaesthesia*. Published electronically February 27, 2019. doi: 10.1016/j.bja.2019.01.013

2026 AQR QCDR Measure Specifications Manual

Sesso, Howard D., Meir J. Stampfer, Bernard Rosner, Charles H. Hennekens, J. Michael Gaziano, JoAnn E. Manson, and Robert J. Glynn. "Systolic and Diastolic Blood Pressure, Pulse Pressure, and Mean Arterial Pressure as Predictors of Cardiovascular Disease Risk in Men." *Hypertension*, vol. 36, no. 5, 2000, pp. 801–807. doi: 10.1161/01.HYP.36.5.801.

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI18: Coronary Artery Bypass Graft (CABG): Prolonged Intubation – Inverse Measure

Provision licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description:

Percentage of patients aged 18 years and older undergoing isolated CABG surgery who require postoperative intubation greater than 24 hours.

Measure Type

Outcome

High Priority Status

Yes

Inverse Measure

Yes

Instructions

This measure is to be reported each time an isolated CABG procedure is performed during the reporting period. It is anticipated that qualified anesthesia providers and eligible clinicians who provide services for isolated CABG will submit this measure. This measure is intended to reflect the quality of services provided for isolated CABG or isolated reoperation CABG patients.

Measure Reporting via the Qualified Clinical Data Registry

CPT codes and patient demographics are used to identify patients who are included in the measure's denominator. The measure must capture both the surgical and related anesthesia code. G-codes are used to report the numerator of the measure.

Denominator

All patients, aged 18 years and older, undergoing isolated CABG surgery

Definition: Isolated CABG refers to CABG using arterial and/or venous grafts only.

Denominator Criteria (Eligible Cases):

Patient aged 18 years and older on date of encounter

AND

Patient encounter during the reporting period (CPT): 33510, 33511, 33512, 33513, 33514, 33516, 33517, 33518, 33519, 33521, 33522, 33523, 33533, 33534, 33535, 33536

AND

00566, 00567

OR

Patient encounter during the reporting period (CPT): 33510, 33511, 33512, 33513, 33514, 33516, 33517, 33518, 33519, 33521, 33522, 33523, 33533, 33534, 33535, 33536

AND

Patient encounter during the reporting period (CPT): 33530

AND

00562

2026 AQR QCDR Measure Specifications Manual

Denominator Exclusions

- Organ donors as designated by ASA Physical Status 6
- Procedure reduced or discontinued prior to initiation of CPB as indicated on the claim by Modifier 52 or Modifier 53
- Transcatheter aortic valve replacement (TAVR) or transcatheter aortic valve implantation (TAVI) cases (CPT codes 33361, 33362, 33363, 33364, 33365, 33366, 33367, 33368, 33369, 93355): **12A38**

Numerator

Patients who require intubation greater than 24 hours following exit from the operating room

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

G8569 Prolonged postoperative intubation (> 24 hrs) required

OR

Performance Not Met:

G8570 Prolonged postoperative intubation (>24 hrs) not required

CBE Number: Not applicable

eCQM: Not applicable

Data Source: Hybrid; Other: Other: Medical Record, Registry

Care Setting: Hospital Inpatient

Telehealth: No

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number of Performance Rates: 1

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI48: Patient-Reported Experience with Anesthesia

Provision licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description: Percentage of patients, aged 18 and older, who were surveyed on their patient experience and satisfaction with anesthesia care and who reported a positive experience.

This measure will consist of two performance rates:

AQI48a: Percentage of patients, aged 18 and older, who were surveyed on their patient experience and satisfaction with anesthesia care

AQI48b: Percentage of patients, aged 18 and older, who completed a survey on their patient experience and satisfaction with anesthesia care who report a positive experience with anesthesia care within 60 days of receipt of the survey.

*NOTE: The measure requires that a valid survey, as defined in the numerator of AQI48a, be sent to patients between discharge from the facility and within 30 days of facility discharge. To report AQI48b, a minimum number of 20 surveys with the mandatory question completed must be reported. ** In order to be scored on this measure, clinicians must report BOTH AQI48a AND AQI48b.*

Measure Type

Patient-Reported Outcome-based Performance Measure (PRO-PM)

High Priority Status

Yes

Inverse Measure:

No

Instructions:

This measure, consisting of two performance rates for AQI48a and AQI48b, is to be reported each time a patient underwent a procedure* with anesthesia during the reporting period. AQI48a should be reported each time a patient undergoes a procedure under anesthesia. To report AQI48b, the provider must report the individual patient scores received by the patient who completed the survey described in AQI48a. A percentage reporting a positive experience will be calculated by the registry on the provider's behalf. It is anticipated that qualified anesthesia providers and eligible clinicians who provide denominator-eligible services will submit this measure.

Measure Reporting via the Qualified Clinical Data Registry

CPT codes and patient demographics are used to identify patients who are included in the measure denominator. Registry codes are used to report the measure numerator.

Denominator

Patients, aged 18 and older, who undergo a procedure* under anesthesia (AQI48a) and who complete a survey on their patient experience and satisfaction with anesthesia care within 60 days of procedure (AQI48b)

Definition: *Any procedure including surgical, therapeutic or diagnostic

Denominator Note: In order to report AQI48b, the denominator must include a minimum of 20 returned surveys.

2026 AQR QCDR Measure Specifications Manual

Denominator-AQI48a

Patients aged 18 and older, who undergo a procedure* under anesthesia
Definition: *Any procedure including surgical, therapeutic or diagnostic

Denominator-AQI48b

All patients from the numerator of AQI48a who complete a survey on their patient experience and satisfaction with anesthesia care within 60 days of receipt of the survey.

Denominator Criteria (Eligible Cases):

Patient aged 18 years or older on date of encounter

AND

AQI 48a: Patient encounter during the reporting period (CPT):

00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00162, 00164, 00170, 00172, 00174, 00176, 00190, 00192, 00210, 00211, 00212, 00214, 00215, 00216, 00218, 00220, 00222, 00300, 00320, 00322, 00350, 00352, 00400, 00402, 00404, 00406, 00410, 00450, 00454, 00470, 00472, 00474, 00500, 00520, 00522, 00524, 00528, 00529, 00530, 00532, 00534, 00537, 00539, 00540, 00541, 00542, 00546, 00548, 00550, 00560, 00562, 00563, 00566, 00567, 00580, 00600, 00604, 00620, 00625, 00626, 00630, 00632, 00635, 00640, 00670, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00754, 00756, 00770, 00790, 00792, 00794, 00796, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00840, 00842, 00844, 00846, 00848, 00851, 00860, 00862, 00864, 00865, 00866, 00868, 00870, 00872, 00873, 00880, 00882, 00902, 00904, 00906, 00908, 00910, 00912, 00914, 00916, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00932, 00934, 00936, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01130, 01140, 01150, 01160, 01170, 01173, 01200, 01202, 01210, 01212, 01214, 01215, 01220, 01230, 01232, 01234, 01250, 01260, 01270, 01272, 01274, 01320, 01340, 01360, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01420, 01430, 01432, 01440, 01442, 01444, 01462, 01464, 01470, 01472, 01474, 01480, 01482, 01484, 01486, 01490, 01500, 01502, 01520, 01522, 01610, 01620, 01622, 01630, 01634, 01636, 01638, 01650, 01652, 01654, 01656, 01670, 01680, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01756, 01758, 01760, 01770, 01772, 01780, 01782, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920, 01922, 01924, 01925, 01926, 01930, 01931, 01932, 01933, 01937, 01938, 01939, 01940, 01941, 01942, 01951, 01952, 01958, 01960, 01961, 01962, 01963, 01965, 01966, 01967, 20526, 20550, 20551, 20552, 20553, 20600, 20604, 20605, 20606, 20610, 20611, 27096, 36555, 36556, 36570, 36571, 36578, 36580, 36581, 36582, 36583, 36584, 36585, 62263, 62264, 62270, 62272, 62273, 62280, 62281, 62282, 62320, 62321, 62322, 62323, 62324, 62325, 62326, 62327, 62328, 62329, 62350, 62355, 62360, 62361, 62362, 62365, 62370, 63650, 63661, 63662, 63663, 63664, 63685, 63688, 64400, 64405, 64408, 64415, 64416, 64417, 64418, 64420, 64425, 64430, 64435, 64445, 64446, 64447, 64448, 64449, 64450, 64451, 64454, 64461, 64463, 64479, 64483, 64486, 64487, 64488, 64489, 64490, 64493, 64505, 64510, 64517, 64520, 64530, 64600, 64605, 64610, 64620, 64624, 64625, 64630, 64633, 64635, 64640, 64680, 64681, 93503, 95990, 95991

For AQI48b

AND

Patient completed a survey on their patient experience and satisfaction with anesthesia care within 60 days of receipt of the survey **11A94**

Denominator Exclusions

- **48a:** Organ Donors as designated with ASA Physical Status 6
- **48a:** Patient died within 30 days of the procedure: **10A11**
- **48b:** Patient did not complete the mandatory anesthesia satisfaction question: **11A95**

Numerator-AQI48a:

Patients who received a survey within 30 days of the procedure to assess their experience and satisfaction with anesthesia.

Numerator Note: The survey should be administered to the patient shortly following discharge from the facility.

2026 AQR QCDR Measure Specifications Manual

Definition: Practices and eligible clinicians may customize their patient experience and satisfaction with anesthesia surveys to meet local needs but, at a minimum, a valid survey must include a core set of questions that address three of the four following criteria related to patient experience and satisfaction and one mandatory question described below.

1. Pre-operative Education and Preparation
2. Patient and/or Family Communication
3. Care Team Response to Comfort and Well-Being
4. Post-operative pain control and/or management

Mandatory question that must be included in each valid survey (practices must also include an option for patient to indicate “Not Applicable”):

1. On a scale of 1 to 5, where 1 indicates the worst anesthesia experience and where 5 indicates the best anesthesia experience, how would you rate your anesthesia experience?

Numerator Note: Practices and eligible clinicians may wish to supplement these questions by taking into consideration the recommendations of the ASA Committee on Performance and Outcomes Measurement work product entitled “Patient Satisfaction and Experience with Anesthesia.”

Numerator Note: Depending on local practice, practices and eligible clinicians may wish to supplement survey questions by taking into consideration the recommendations developed as part of the Perioperative Surgical Home (PSH) that are structured in five distinct components.

1. Pre-Operative Education and Preparation (Four Indicators)
 - a. Patient comfort with instructions provided about eating better
 - b. Patient comfort with instructions provided about exercise or physical therapy
 - c. Patient comfort with instructions provided about stopping smoking (if applicable)
 - d. Patient comfort with instructions provided about what to do after surgery
2. Check-In and Pre-Procedure Experience
3. Caregiver and Family Communication during Surgery
4. Care Team Response to Comfort and Well-Being
5. Post-Operative Pain Management

For more information on these resources, visit <https://www.asahq.org/psh>.

Numerator Quality-Data Coding Options for Reporting Satisfactorily: AQI48a

Performance Met:

10A12 Patient provided with a survey within 30 days of the procedure to assess their experience and satisfaction with anesthesia

OR

Denominator Exception

10A13 Documentation of patient reason(s), process reason(s) or medical reason(s) for not receiving survey (i.e. patients who are non-verbal, who are unable to be surveyed due to a medical or psychiatric reason, who are unable to be surveyed due to a language barrier, have not provided contact information, who are discharged to assisted living, skilled nursing facility or other similar location where direct access to the patient is not available, or who decline to be surveyed)

OR

Performance Not Met:

10A14 Patient was not provided with a survey within 30 days of the procedure to assess their experience and satisfaction with anesthesia

2026 AQR QCDR Measure Specifications Manual

OR

11A93 The patient did not respond within 60 days of receipt of the survey. *

*The intent of this registry code is to eliminate the surveys that were not returned within the required 60-day period. Reporting this code is essential for the data completeness calculation and does not indicate negative performance of this measure. Performance is calculated based on performance of AQI48b.

Numerator- AQI 48b:

Patients who reported a positive experience with anesthesia care within 60 days of receipt of the survey.

Definition: A positive experience is defined as a response of 4 or 5 on the following mandatory patient experience and satisfaction survey question:

On a scale of 1 to 5, where 1 indicates the worst anesthesia experience and where 5 indicates the best anesthesia experience, how would you rate your overall anesthesia experience? (*Practices must include an option for patient to indicate "Not Applicable"*)

Numerator Quality-Data Coding Options for Reporting Satisfactorily: AQI48b

Reporting note: To report this measure, the provider must report the individual patient scores. A percentage reporting a positive experience will be calculated on the provider's behalf.

Performance Met:

10A70 Patient reported a positive anesthesia experience (i.e., a 4 or 5 on the mandatory survey question)

OR

Performance Not Met:

10A71 Patient did NOT report a positive anesthesia experience (i.e., a 1, 2, or 3 on the mandatory survey question)

Data Source: Hybrid; Other: Other: Quality measures included in AQI NACOR are reported to the registry by participants who use a combination data sources that may include one or more of the following: administrative claims/billing data, facility discharge data, EHR (AIMS, partial patient record), paper medical record, and/or contracted third party data capture systems

Care Settings: Ambulatory Care: Clinician Office/Clinic; Ambulatory Care: Hospital; Hospital; Hospital Inpatient; Outpatient Services

Telehealth: No

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number of Multiple Performance Rates: 2

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Ratio Measure: No

Risk Adjusted: No

MIPS Reporting Option: Traditional MIPS, MVP

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI65: Avoidance of Cerebral Hyperthermia for Procedures Involving Cardiopulmonary Bypass

Provision licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description: Percentage of patients, aged 18 years and older, undergoing a procedure using cardiopulmonary bypass who did not have a documented intraoperative pulmonary artery, oropharyngeal, or nasopharyngeal temperature ≥ 37.0 degrees Celsius during the period of cardiopulmonary bypass

Measure Type

Outcome

High Priority Status

Yes

Inverse Measure

No

Instructions

This measure is to be reported each time a patient undergoes a cardiac operation using cardiopulmonary bypass during the reporting period. It is anticipated that qualified anesthesia providers and eligible clinicians who provide denominator-eligible services will submit this measure.

Measure Reporting via the Qualified Clinical Data Registry

CPT codes and patient demographics are used to identify patients who are included in the measure's denominator. Registry codes are used to report the numerator.

Denominator

All patients aged 18 years or older, who undergo a procedure using cardiopulmonary bypass

Denominator Criteria (Eligible Cases):

Patient aged 18 years and older

AND

Patient encounter during the reporting period (CPT): 00562, 00563, 00567, 00580

Denominator Exclusions

- Procedure reduced or discontinued prior to initiation of CPB as indicated on the claim by Modifier 52 or Modifier 53.
- Transcatheter aortic valve replacement (TAVR) or transcatheter aortic valve implantation (TAVI) cases (CPT codes 33361, 33362, 33363, 33364, 33365, 33366, 33367, 33368, 33369, 93355): **12A38**

Numerator

Patients who did not have an intraoperative pulmonary artery, oropharyngeal, or nasopharyngeal temperature ≥ 37.0 degrees Celsius during cardiopulmonary bypass

2026 AQR QCDR Measure Specifications Manual

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

11A11 All intraoperative pulmonary artery, oropharyngeal, or nasopharyngeal temperatures <37.0 degrees Celsius during cardiopulmonary bypass

OR

Performance Not Met:

11A12 At least one intraoperative pulmonary artery, oropharyngeal, or nasopharyngeal temperature \geq 37.0 degrees Celsius

OR

11A13 No documented pulmonary artery, oropharyngeal, or nasopharyngeal temperatures during cardiopulmonary bypass

CBE Number: Not applicable

eCQM: Not applicable

Data Source: Hybrid; Other: Other: Quality measures included in AQI NACOR are reported to the registry by participants who use a combination data sources that may include one or more of the following: administrative claims/billing data, facility discharge data, EHR (AIMS, partial patient record), paper medical record, and/or contracted third party data capture systems

Care Setting: Hospital Inpatient

Telehealth: No

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number of Multiple Performance Rates: Not applicable

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Ratio Measure: No

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI71: Ambulatory Glucose Management

Provision licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description: Percentage of diabetic patients, aged 18 years and older, who receive an office-based or ambulatory surgery whose blood glucose level is appropriately managed throughout the perioperative period.

This measure will consist of four performance rates:

AQI71a: Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery whose blood glucose level is tested prior to the start of anesthesia.

AQI71b: Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L) who received insulin prior to anesthesia end time.

AQI71c: Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who received insulin perioperatively and who received a follow-up blood glucose level check following the administration of insulin and prior to discharge.

AQI71d: Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L) who received education on managing their glucose in the postoperative period prior to discharge.

NOTE: The overall measure score will be calculated as an average of the performance rates of parts A, B, C and D. In order to be scored on this measure, clinicians must have at least one eligible case reported for each sub-metric: AQI71a, AQI71b, AQI71c, and AQI71d.

Measure Type

Composite

High Priority Status

Yes

Inverse Measure

No

Instructions

This measure will consist of four performance rates: AQI71a, AQI71b, AQI71c, and AQI71d. Each measure should be reported, as appropriate, for each time a patient undergoes a procedure in an office-based or ambulatory setting during the reporting period. This measure has four sub-metrics which are used to calculate the total composite score. All sub-metrics are required to be reported during the performance period. In order to be scored on this measure, clinicians must have at least one eligible case reported for AQI71a, AQI71b, AQI71c, and AQI71d. It is anticipated that qualified anesthesia providers and eligible clinicians who provide denominator-eligible services will submit this measure.

Measure Reporting via the Qualified Clinical Data Registry

CPT codes and patient demographics are used to identify patients who are included in the measure's denominator.

Registry codes are used to report the numerator of the measure.

2026 AQR QCDR Measure Specifications Manual

CBE Number: Not Applicable

eCQM: Not Applicable

Data Source: Hybrid; Other: Other: We expect that the data source used for the measure will be similar to other AQI NACOR QCDR Measures. Quality measures included in AQI NACOR are reported to the registry by participants who use a combination data sources that may include one or more of the following: administrative claims/billing data, facility discharge data, EHR (AIMS, partial patient record), paper medical record, and/or contracted third party data capture systems.

Care Setting: Ambulatory Care: Hospital

Telehealth: No

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number of Performance Rates: 5

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS

AQI71a: Ambulatory Point-of-Care Glucose Testing

Description: Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery whose blood glucose level is tested prior to the start of anesthesia.

Denominator:

All patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery

Denominator definition: Office-based and ambulatory surgery is defined as a therapeutic or diagnostic procedure performed in a healthcare facility that does not require an overnight stay (less than 24 hours of care)

Denominator Criteria (eligible cases):

All patients, aged 18 years and older

AND

Diagnosis of diabetes mellitus: **11A41**

OR

ICD-10CM code: E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.3211, E10.3212, E10.3213, E10.3219, E10.3291, E10.3292, E10.3293, E10.3299, E10.3311, E10.3312, E10.3313, E10.3319, E10.3391, E10.3392, E10.3393, E10.3399, E10.3411, E10.3412, E10.3413, E10.3419, E10.3491, E10.3492, E10.3493, E10.3499, E10.3511, E10.3512, E10.3513, E10.3519, E10.3521, E10.3522, E10.3523, E10.3529, E10.3531, E10.3532, E10.3533, E10.3539, E10.3541,

2026 AQR QCDR Measure Specifications Manual

E10.3542, E10.3543, E10.3549, E10.3551, E10.3552, E10.3553, E10.3559, E10.3591, E10.3592, E10.3593, E10.3599, E10.36, E10.37X1, E10.37X2, E10.37X3, E10.37X9, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9, E10.A0, E10.A1, E10.A2, E11.00, E11.01, E11.10, E11.11, E11.21, E11.22, E11.29, E11.311, E11.319, E11.3211, E11.3212, E11.3213, E11.3219, E11.3291, E11.3292, E11.3293, E11.3299, E11.3311, E11.3312, E11.3313, E11.3319, E11.3391, E11.3392, E11.3393, E11.3399, E11.3411, E11.3412, E11.3413, E11.3419, E11.3491, E11.3492, E11.3493, E11.3499, E11.3511, E11.3512, E11.3513, E11.3519, E11.3521, E11.3522, E11.3523, E11.3529, E11.3531, E11.3532, E11.3533, E11.3539, E11.3541, E11.3542, E11.3543, E11.3549, E11.3551, E11.3552, E11.3553, E11.3559, E11.3591, E11.3592, E11.3593, E11.3599, E11.36, E11.37X1, E11.37X2, E11.37X3, E11.37X9, E11.39, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9, E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.3211, E13.3212, E13.3213, E13.3219, E13.3291, E13.3292, E13.3293, E13.3299, E13.3311, E13.3312, E13.3313, E13.3319, E13.3391, E13.3392, E13.3393, E13.3399, E13.3411, E13.3412, E13.3413, E13.3419, E13.3491, E13.3492, E13.3493, E13.3499, E13.3511, E13.3512, E13.3513, E13.3519, E13.3521, E13.3522, E13.3523, E13.3529, E13.3531, E13.3532, E13.3533, E13.3539, E13.3541, E13.3542, E13.3543, E13.3549, E13.3551, E13.3552, E13.3553, E13.3559, E13.3591, E13.3592, E13.3593, E13.3599, E13.36, E13.37X1, E13.37X2, E13.37X3, E13.37X9, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9

AND

Off Campus-Outpatient Hospital, On Campus – Outpatient Hospital, Ambulatory Surgical Center, and Office-Based Setting: Place of Service Codes 11, 19, 22 or 24

AND

Patient encounter during the reporting period (CPT): 00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00164, 00170, 00172, 00174, 00176, 00300, 00320, 00322, 00400, 00402, 00404, 00410, 00450, 00454, 00520, 00522, 00524, 00530, 00532, 00534, 00537, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00790, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00851, 00870, 00872, 00873, 00902, 00906, 00910, 00912, 00914, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01202, 01250, 01260, 01320, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01464, 01470, 01472, 01474, 01480, 01520, 01610, 01620, 01622, 01630, 01634, 01638, 01670, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01758, 01760, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920, 01930, 01937, 01938, 01939, 01940, 01941, 01942, 01965, 01966

Denominator Exclusions:

- Procedure <30 minutes duration: **11A45**

Numerator:

Patients who received a blood glucose test prior to the start of anesthesia

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

11A51 Patient received a blood glucose test prior to start of anesthesia

OR

Performance Not Met:

11A52 Patient did NOT receive a glucose test prior to start of anesthesia

2026 AQR QCDR Measure Specifications Manual

AQI71b: Ambulatory Hyperglycemia Control

Description

Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L) who received insulin prior to anesthesia end time.

Denominator:

All patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L)

Denominator definition: Office-based and ambulatory surgery is defined as a therapeutic or diagnostic procedure performed in a healthcare facility that does not require an overnight stay (less than 24 hours of care)

Denominator Criteria (Eligible Cases):

All patients, aged 18 years and older

AND

Diagnosis of diabetes mellitus: **11A41**

OR

ICD-10CM code: E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.3211, E10.3212, E10.3213, E10.3219, E10.3291, E10.3292, E10.3293, E10.3299, E10.3311, E10.3312, E10.3313, E10.3319, E10.3391, E10.3392, E10.3393, E10.3399, E10.3411, E10.3412, E10.3413, E10.3419, E10.3491, E10.3492, E10.3493, E10.3499, E10.3511, E10.3512, E10.3513, E10.3519, E10.3521, E10.3522, E10.3523, E10.3529, E10.3531, E10.3532, E10.3533, E10.3539, E10.3541, E10.3542, E10.3543, E10.3549, E10.3551, E10.3552, E10.3553, E10.3559, E10.3591, E10.3592, E10.3593, E10.3599, E10.36, E10.37X1, E10.37X2, E10.37X3, E10.37X9, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9, E10.A0, E10.A1, E10.A2, E11.00, E11.01, E11.10, E11.11, E11.21, E11.22, E11.29, E11.311, E11.319, E11.3211, E11.3212, E11.3213, E11.3219, E11.3291, E11.3292, E11.3293, E11.3299, E11.3311, E11.3312, E11.3313, E11.3319, E11.3391, E11.3392, E11.3393, E11.3399, E11.3411, E11.3412, E11.3413, E11.3419, E11.3491, E11.3492, E11.3493, E11.3499, E11.3511, E11.3512, E11.3513, E11.3519, E11.3521, E11.3522, E11.3523, E11.3529, E11.3531, E11.3532, E11.3533, E11.3539, E11.3541, E11.3542, E11.3543, E11.3549, E11.3551, E11.3552, E11.3553, E11.3559, E11.3591, E11.3592, E11.3593, E11.3599, E11.36, E11.37X1, E11.37X2, E11.37X3, E11.37X9, E11.39, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9, E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.3211, E13.3212, E13.3213, E13.3219, E13.3291, E13.3292, E13.3293, E13.3299, E13.3311, E13.3312, E13.3313, E13.3319, E13.3391, E13.3392, E13.3393, E13.3399, E13.3411, E13.3412, E13.3413, E13.3419, E13.3491, E13.3492, E13.3493, E13.3499, E13.3511, E13.3512, E13.3513, E13.3519, E13.3521, E13.3522, E13.3523, E13.3529, E13.3531, E13.3532, E13.3533, E13.3539, E13.3541, E13.3542, E13.3543, E13.3549, E13.3551, E13.3552, E13.3553, E13.3559, E13.3591, E13.3592, E13.3593, E13.3599, E13.36, E13.37X1, E13.37X2, E13.37X3, E13.37X9, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9

AND

Off Campus-Outpatient Hospital, On Campus – Outpatient Hospital, Ambulatory Surgical Center, and Office-Based Setting: Place of Service Codes 11, 19, 22 or 24

AND

Experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L) prior to anesthesia end time: **11A44**

AND

2026 AQR QCDR Measure Specifications Manual

Patient encounter during the reporting period (CPT): 00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00164, 00170, 00172, 00174, 00176, 00300, 00320, 00322, 00400, 00402, 00404, 00410, 00450, 00454, 00520, 00522, 00524, 00530, 00532, 00534, 00537, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00790, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00851, 00870, 00872, 00873, 00902, 00906, 00910, 00912, 00914, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01202, 01250, 01260, 01320, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01464, 01470, 01472, 01474, 01480, 01520, 01610, 01620, 01622, 01630, 01634, 01638, 01670, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01758, 01760, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920, 01930, 01937, 01938, 01939, 01940, 01941, 01942, 01965, 01966

Denominator Exclusions:

- Procedure <30 minutes duration: **11A45**

Numerator:

Patients who received insulin prior to anesthesia end time.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

11A53 Patient received insulin prior to anesthesia end time.

OR

Denominator Exception:

11A82 Documentation that insulin was not given because patient had severe comorbidities and glucose concentrations between 180 mg/dL and 250 mg/dL (10-13.9 mmol/L).

OR

Performance Not Met:

11A54 Patient did NOT receive insulin prior to anesthesia end time.

AQI71c: Follow-Up Glucose Check for Patients Receiving Insulin

Description: Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who received insulin perioperatively and who received a follow-up blood glucose level check following the administration of insulin and prior to discharge.

Denominator:

All patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who received insulin perioperatively

Denominator definition: Office-based and ambulatory surgery is defined as a therapeutic or diagnostic procedure performed in a healthcare facility that does not require an overnight stay (less than 24 hours of care)

Denominator Criteria (Eligible Cases):

All patients, aged 18 years and older

AND

Diagnosis of diabetes mellitus: **11A41**

OR

ICD-10CM code: E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.3211, E10.3212, E10.3213, E10.3219, E10.3291, E10.3292, E10.3293, E10.3299, E10.3311, E10.3312, E10.3313, E10.3319, E10.3391, E10.3392, E10.3393, E10.3399, E10.3411, E10.3412, E10.3413, E10.3419, E10.3491, E10.3492, E10.3493, E10.3499, E10.3511, E10.3512,

2026 AQR QCDR Measure Specifications Manual

E10.3513, E10.3519, E10.3521, E10.3522, E10.3523, E10.3529, E10.3531, E10.3532, E10.3533, E10.3539, E10.3541, E10.3542, E10.3543, E10.3549, E10.3551, E10.3552, E10.3553, E10.3559, E10.3591, E10.3592, E10.3593, E10.3599, E10.36, E10.37X1, E10.37X2, E10.37X3, E10.37X9, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9, E10.A0, E10.A1, E10.A2, E11.00, E11.01, E11.10, E11.11, E11.21, E11.22, E11.29, E11.311, E11.319, E11.3211, E11.3212, E11.3213, E11.3219, E11.3291, E11.3292, E11.3293, E11.3299, E11.3311, E11.3312, E11.3313, E11.3319, E11.3391, E11.3392, E11.3393, E11.3399, E11.3411, E11.3412, E11.3413, E11.3419, E11.3491, E11.3492, E11.3493, E11.3499, E11.3511, E11.3512, E11.3513, E11.3519, E11.3521, E11.3522, E11.3523, E11.3529, E11.3531, E11.3532, E11.3533, E11.3539, E11.3541, E11.3542, E11.3543, E11.3549, E11.3551, E11.3552, E11.3553, E11.3559, E11.3591, E11.3592, E11.3593, E11.3599, E11.36, E11.37X1, E11.37X2, E11.37X3, E11.37X9, E11.39, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9, E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.3211, E13.3212, E13.3213, E13.3219, E13.3291, E13.3292, E13.3293, E13.3299, E13.3311, E13.3312, E13.3313, E13.3319, E13.3391, E13.3392, E13.3393, E13.3399, E13.3411, E13.3412, E13.3413, E13.3419, E13.3491, E13.3492, E13.3493, E13.3499, E13.3511, E13.3512, E13.3513, E13.3519, E13.3521, E13.3522, E13.3523, E13.3529, E13.3531, E13.3532, E13.3533, E13.3539, E13.3541, E13.3542, E13.3543, E13.3549, E13.3551, E13.3552, E13.3553, E13.3559, E13.3591, E13.3592, E13.3593, E13.3599, E13.36, E13.37X1, E13.37X2, E13.37X3, E13.37X9, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9

AND

Off Campus-Outpatient Hospital, On Campus – Outpatient Hospital, Ambulatory Surgical Center, and Office-Based Setting: Place of Service Codes 11, 19, 22 or 24

AND

Patient received insulin perioperatively: **11A55**

AND

Patient encounter during the reporting period (CPT): 00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00164, 00170, 00172, 00174, 00176, 00300, 00320, 00322, 00400, 00402, 00404, 00410, 00450, 00454, 00520, 00522, 00524, 00530, 00532, 00534, 00537, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00790, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00851, 00870, 00872, 00873, 00902, 00906, 00910, 00912, 00914, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01202, 01250, 01260, 01320, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01464, 01470, 01472, 01474, 01480, 01520, 01610, 01620, 01622, 01630, 01634, 01638, 01670, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01758, 01760, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920, 01930, 01937, 01938, 01939, 01940, 01941, 01942, 01965, 01966

Denominator Exclusions:

- Procedure <30 minutes duration: **11A45**

Numerator:

Patients who received a follow-up blood glucose level check following the administration of insulin and prior to discharge.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

11A56 Patient received a follow-up blood glucose level check following the administration of insulin and prior to discharge.

OR

2026 AQR QCDR Measure Specifications Manual

Performance Not Met:

11A57 Patient did NOT receive a follow-up blood glucose level check following the administration of insulin and prior to discharge.

AQI71d: Hyperglycemia Management Patient Education

Description: Percentage of patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L) who received education on managing their glucose in the postoperative period prior to discharge.

Denominator:

All patients, aged 18 years and older, with a current diagnosis of diabetes mellitus receiving anesthesia services for office-based or ambulatory surgery who experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L).

Denominator definition: Office-based or ambulatory surgery is defined as a therapeutic or diagnostic procedure performed in a healthcare facility that does not require an overnight stay (less than 24 hours of care).

Denominator Criteria (Eligible Cases):

All patients, aged 18 years and older

AND

Diagnosis of diabetes mellitus: **11A41**

OR

ICD-10CM code: E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.3211, E10.3212, E10.3213, E10.3219, E10.3291, E10.3292, E10.3293, E10.3299, E10.3311, E10.3312, E10.3313, E10.3319, E10.3391, E10.3392, E10.3393, E10.3399, E10.3411, E10.3412, E10.3413, E10.3419, E10.3491, E10.3492, E10.3493, E10.3499, E10.3511, E10.3512, E10.3513, E10.3519, E10.3521, E10.3522, E10.3523, E10.3529, E10.3531, E10.3532, E10.3533, E10.3539, E10.3541, E10.3542, E10.3543, E10.3549, E10.3551, E10.3552, E10.3553, E10.3559, E10.3591, E10.3592, E10.3593, E10.3599, E10.36, E10.37X1, E10.37X2, E10.37X3, E10.37X9, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9, E10.A0, E10.A1, E10.A2, E11.00, E11.01, E11.10, E11.11, E11.21, E11.22, E11.29, E11.311, E11.319, E11.3211, E11.3212, E11.3213, E11.3219, E11.3291, E11.3292, E11.3293, E11.3299, E11.3311, E11.3312, E11.3313, E11.3319, E11.3391, E11.3392, E11.3393, E11.3399, E11.3411, E11.3412, E11.3413, E11.3419, E11.3491, E11.3492, E11.3493, E11.3499, E11.3511, E11.3512, E11.3513, E11.3519, E11.3521, E11.3522, E11.3523, E11.3529, E11.3531, E11.3532, E11.3533, E11.3539, E11.3541, E11.3542, E11.3543, E11.3549, E11.3551, E11.3552, E11.3553, E11.3559, E11.3591, E11.3592, E11.3593, E11.3599, E11.36, E11.37X1, E11.37X2, E11.37X3, E11.37X9, E11.39, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9, E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.3211, E13.3212, E13.3213, E13.3219, E13.3291, E13.3292, E13.3293, E13.3299, E13.3311, E13.3312, E13.3313, E13.3319, E13.3391, E13.3392, E13.3393, E13.3399, E13.3411, E13.3412, E13.3413, E13.3419, E13.3491, E13.3492, E13.3493, E13.3499, E13.3511, E13.3512, E13.3513, E13.3519, E13.3521, E13.3522, E13.3523, E13.3529, E13.3531, E13.3532, E13.3533, E13.3539, E13.3541, E13.3542, E13.3543, E13.3549, E13.3551, E13.3552, E13.3553, E13.3559, E13.3591, E13.3592, E13.3593, E13.3599, E13.36, E13.37X1, E13.37X2, E13.37X3, E13.37X9, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9

AND

Off Campus-Outpatient Hospital, On Campus – Outpatient Hospital, Ambulatory Surgical Center, and Office-Based Setting: Place of Service Codes 11, 19, 22 or 24

AND

2026 AQR QCDR Measure Specifications Manual

Experienced a blood glucose level ≥ 180 mg/dL (10.0 mmol/L) prior to anesthesia end time: **11A44**

AND

Patient encounter during the reporting period (CPT): 00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00164, 00170, 00172, 00174, 00176, 00300, 00320, 00322, 00400, 00402, 00404, 00410, 00450, 00454, 00520, 00522, 00524, 00530, 00532, 00534, 00537, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00790, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00851, 00870, 00872, 00873, 00902, 00906, 00910, 00912, 00914, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01202, 01250, 01260, 01320, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01464, 01470, 01472, 01474, 01480, 01520, 01610, 01620, 01622, 01630, 01634, 01638, 01670, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01758, 01760, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920, 01930, 01937, 01938, 01939, 01940, 01941, 01942, 01965, 01966

Denominator Exclusions:

- Procedure <30 minutes duration: **11A45**

Numerator:

Patients who received education on managing their glucose in the postoperative period prior to discharge

Numerator Note: To meet this measure, the anesthesiologist or other member of the care team must provide both oral and written education. Provision of written materials alone is not sufficient.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

11A58 Patient received education on managing their glucose in the postoperative period prior to discharge.

OR

Performance Not Met:

11A59 Patient did NOT receive education on managing their glucose in the postoperative period prior to discharge.

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI79: Neuromuscular Blockade and Quantitative Train-of-Four

Provision licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description

The percentage of patients, 18 years and older, who undergo a procedure utilizing non-depolarizing neuromuscular blockade (NMB) who have a documented quantitative train-of-four ratio of greater than or equal to 0.9 before tracheal extubation in the operating room, post-anesthesia care unit, non-operating room anesthetizing location or earliest extubation.

Measure Type

Process

High Priority Status

Yes

Inverse Measure

No

Instructions

This measure is to be submitted each time a patient undergoes a surgical, therapeutic, or diagnostic procedure using a non-depolarizing neuromuscular blockade medication is performed during the performance period. There is no diagnosis associated with this measure. It is anticipated that Merit-based Incentive Payment System (MIPS) eligible clinicians who provide the listed anesthesia services as specified in the denominator coding will submit this measure.

Measure Reporting via the Qualified Clinical Data Registry

CPT (Current Procedural Terminology) codes, denominator criteria, registry codes, and patient demographics are used to identify patients who are included in the measure's denominator. The numerator options included in this specification are used to submit the quality actions as allowed by the measure.

Denominator

All patients, aged 1 and above, who undergo a surgical, therapeutic, or diagnostic procedure that involves using a non-depolarizing neuromuscular blockade (NMB).

Denominator Note: For this measure, qualifying neuromuscular blocker medications include:

- Atracurium
- Cisatracurium
- Pancuronium
- Rocuronium
- Vecuronium

Denominator Criteria (Eligible Cases):

All patients aged 1 year and older

AND

Patient encounter during the reporting period (CPT):

00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00162, 00164, 00170, 00172, 00174, 00176, 00190, 00192, 00210, 00211, 00212, 00214, 00215, 00216, 00218, 00220, 00222, 00300, 00320, 00322, 00350, 00352, 00400, 00402, 00404, 00406, 00410, 00450, 00454, 00470, 00472, 00474, 00500, 00520,

2026 AQR QCDR Measure Specifications Manual

00522, 00524, 00528, 00529, 00530, 00532, 00534, 00537, 00539, 00540, 00541, 00542, 00546, 00548, 00550, 00560, 00562, 00563, 00566, 00567, 00580, 00600, 00604, 00620, 00625, 00626, 00630, 00632, 00635, 00640, 00670, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00754, 00756, 00770, 00790, 00792, 00794, 00796, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00840, 00842, 00844, 00846, 00848, 00851, 00860, 00862, 00864, 00865, 00866, 00868, 00870, 00872, 00873, 00880, 00882, 00902, 00904, 00906, 00908, 00910, 00912, 00914, 00916, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00932, 00934, 00936, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01130, 01140, 01150, 01160, 01170, 01173, 01200, 01202, 01210, 01212, 01214, 01215, 01220, 01230, 01232, 01234, 01250, 01260, 01270, 01272, 01274, 01320, 01340, 01360, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01420, 01430, 01432, 01440, 01442, 01444, 01462, 01464, 01470, 01472, 01474, 01480, 01482, 01484, 01486, 01490, 01500, 01502, 01520, 01522, 01610, 01620, 01622, 01630, 01634, 01636, 01638, 01650, 01652, 01654, 01656, 01670, 01680, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01756, 01758, 01760, 01770, 01772, 01780, 01782, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920, 01922, 01924, 01925, 01926, 01930, 01931, 01932, 01933, 01937, 01938, 01939, 01940, 01941, 01942, 01951, 01952, 01958, 01960, 01961, 01962, 01963, 01965, 01966, 01967

AND

Received non-depolarizing neuromuscular blocker (NMB): **11A17**

AND

Tracheal extubation occurred in the operating room, post-anesthesia care unit or non-operating room anesthetizing location or at earliest extubation. **12A31**

Denominator Exclusions

ASA (American Society of Anesthesiologists) Physical Status 5 or 6

Bypass Post Anesthesia Care Unit to Intensive Care Unit. **12A32**

Numerator

Patients with a documented quantitative train-of-four greater than or equal to 0.9 AFTER the last dose or stopping of infusion of neuromuscular blocker and before the patient leaves the operating room, post-anesthesia care unit non-operating room anesthetizing location or earliest tracheal extubation.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

12A39 The patient has a documented quantitative train-of-four of greater than or equal to 0.9 before tracheal extubation in the operating room, post-anesthesia care unit, non-operating room anesthetizing location, or earliest extubation.

OR

Denominator Exception:

12A34 Documentation of the need for continued paralysis postoperatively.

OR

Performance Not Met:

12A40 The patient did NOT have a documented quantitative train-of-four of greater than or equal to 0.9 before tracheal extubation in the operating room, post-anesthesia care unit, non-operating room anesthetizing location, or earliest extubation.

CBE Number: Not applicable

eCQM: Not applicable

Data Source: Administrative claims data; EHR

Care Setting: Ambulatory; Ambulatory Surgical Center; Hospital; Hospital Inpatient; Hospital Outpatient; Office Based Surgery Center

Telehealth: No

2026 AQR QCDR Measure Specifications Manual

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number Performance Rates: 1

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Ratio Measure: No

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI80: Continuation of Buprenorphine for Inpatient Surgical Patients or Methadone Therapy during the Perioperative Period for Inpatient and Ambulatory Surgical Patients

Provision licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description

Percentage of patients, age 13 and older, undergoing an inpatient surgical procedure who have an outpatient prescription for buprenorphine/naloxone or suboxone in their admissions documentation, who are continued on buprenorphine for the duration of their inpatient stay or undergoing an inpatient or ambulatory surgical procedure who are taking methadone as outpatients or who have a prescription for methadone in their admittance documentation, and where methadone is either continued or changed to an equivalent IV dose to maintain the therapeutic level regimen during the perioperative period. For patients receiving care in an ambulatory surgical setting, methadone medication was provided as scheduled before surgery.

NQS Domain / Meaningful Measures Area

Opioid Related

Measure Type

Process

High Priority Status

Yes

Inverse Measure

No

This measure will consist of two performance rates:

1. **Measure A:** Percentage of patients, age 13 and older, undergoing an inpatient surgical procedure who have an outpatient prescription for buprenorphine/naloxone or suboxone in their admissions documentation are continued on buprenorphine for the duration of their inpatient stay.
2. **Measure B:** Percentage of patients, 13 years and older, undergoing an inpatient or ambulatory surgical procedure who are taking methadone as outpatients or who have a prescription for methadone in their admittance documentation and where methadone is either continued or changed to an equivalent IV dose to maintain the therapeutic level regimen during the perioperative period. For patients receiving care in an ambulatory surgical setting, methadone medication was provided as scheduled before surgery.

***NOTE:** The overall measure score will be calculated as an average of the performance rates of A and B. In order to be scored on this measure, clinicians must have at least one eligible case reported for each sub-metric: Measure A and Measure B.*

Measure Instructions:

The MIPS eligible clinician should submit data on one of the submission criteria, depending on the clinical findings and surgical setting. If the patient has an outpatient prescription for buprenorphine/naloxone in their admissions documentation for an inpatient surgical procedure, use **Measure A**.

If the patient is taking methadone as an outpatient, has a prescription for methadone in their admittance documentation, or where methadone medication is provided as scheduled before an ambulatory surgical procedure, use **Measure B**.

2026 AQR QCDR Measure Specifications Manual

There is no diagnosis associated with this measure. It is anticipated that qualified anesthesia professionals and eligible clinicians who provide denominator-eligible services will submit this measure.

Measure Submission Type:

Measure data may be submitted by individual MIPS eligible clinicians, groups, or third-party intermediaries via the Qualified Clinical Data Registry option.

Measure reporting via the Qualified Clinical Data Registry:

CPT (Current Procedural Terminology) Codes, ICD-10 codes, place of service codes, registry codes, denominator criteria, and patient demographics are used to identify patients who are included in the quality performance measure's denominator. The listed denominator criteria are used to identify the intended patient population. The numerator options included in this specification are used to submit the quality actions as allowed by the quality performance measure.

Measure A

All patients aged 13 years and older, undergoing an inpatient surgical procedure, who have an outpatient prescription for buprenorphine/naloxone or suboxone in their admissions documentation

Denominator (Measure A)

All patients aged 18 years or older, who undergo an elective procedure lasting 30 minutes or longer requiring inhalational general anesthesia. (1095)

Denominator Criteria (Eligible Cases):

Patients aged 13 years and older

AND

Patient encounter during the reporting period (CPT):

00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00162, 00164, 00170, 00172, 00174, 00176, 00190, 00192, 00210, 00211, 00212, 00214, 00215, 00216, 00218, 00220, 00222, 00300, 00320, 00322, 00350, 00352, 00400, 00402, 00404, 00406, 00410, 00450, 00454, 00470, 00472, 00474, 00500, 00520, 00522, 00524, 00528, 00529, 00530, 00532, 00534, 00537, 00539, 00540, 00541, 00542, 00546, 00548, 00550, 00560, 00562, 00563, 00566, 00567, 00580, 00600, 00604, 00620, 00625, 00626, 00630, 00632, 00635, 00640, 00670, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00754, 00756, 00770, 00790, 00792, 00794, 00796, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00840, 00842, 00844, 00846, 00848, 00851, 00860, 00862, 00864, 00865, 00866, 00868, 00870, 00872, 00873, 00880, 00882, 00902, 00904, 00906, 00908, 00910, 00912, 00914, 00916, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00932, 00934, 00936, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01130, 01140, 01150, 01160, 01170, 01173, 01200, 01202, 01210, 01212, 01214, 01215, 01220, 01230, 01232, 01234, 01250, 01260, 01270, 01272, 01274, 01320, 01340, 01360, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01420, 01430, 01432, 01440, 01442, 01444, 01462, 01464, 01470, 01472, 01474, 01480, 01482, 01484, 01486, 01490, 01500, 01502, 01520, 01522, 01610, 01620, 01622, 01630, 01634, 01636, 01638, 01650, 01652, 01654, 01656, 01670, 01680, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01756, 01758, 01760, 01770, 01772, 01780, 01782, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920, 01922, 01924, 01925, 01926, 01930, 01931, 01932, 01933, 01937, 01938, 01939, 01940, 01941, 01942, 01951, 01952, 01958, 01960, 01961, 01962, 01963, 01965, 01966, 01967, 01991, 01992

AND

Place of service code: 21

AND

Patient has a prescription for buprenorphine/naloxone or suboxone in their admitting documentation. **12A17**

Denominator Exclusions: Patient has a chronic pain diagnosis (ICD: **G89.4** or Registry Code **12A18**)

2026 AQR QCDR Measure Specifications Manual

Numerator (Measure A)

All patients who receive a continuation of Buprenorphine during their inpatient stay.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

12A19 The patient received a continuation of buprenorphine during their inpatient stay.

OR

Denominator Exception:

12A20 Documentation that the inpatient is unable to tolerate buprenorphine, received instruction from their surgical, anesthesia, and/or addiction treatment provider to stop buprenorphine prior to the inpatient procedure, patient refused continuation of buprenorphine treatment, or received methadone in lieu of buprenorphine.

Performance Not Met:

12A21 The patient did NOT receive a continuation of buprenorphine during their inpatient stay.

Measure B:

Percentage of patients, 13 years and older, undergoing an inpatient or ambulatory surgical procedure who are taking methadone as outpatients or who have a prescription for methadone in their admittance documentation and where methadone is either continued or changed to an equivalent IV dose to maintain the therapeutic level regimen during the perioperative period. For patients receiving care in an ambulatory surgical setting, methadone medication was provided as scheduled before surgery.

Denominator (Measure B):

Patients, 13 years and older, undergoing an inpatient or ambulatory surgical procedure who are taking methadone as outpatients or who have a prescription for methadone in their admittance or ambulatory surgical documentation

Denominator Criteria (Eligible Cases):

Patients aged 13 years or older

AND

Patient encounter during the reporting period (CPT):

00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00162, 00164, 00170, 00172, 00174, 00176, 00190, 00192, 00210, 00211, 00212, 00214, 00215, 00216, 00218, 00220, 00222, 00300, 00320, 00322, 00350, 00352, 00400, 00402, 00404, 00406, 00410, 00450, 00454, 00470, 00472, 00474, 00500, 00520, 00522, 00524, 00528, 00529, 00530, 00532, 00534, 00537, 00539, 00540, 00541, 00542, 00546, 00548, 00550, 00560, 00562, 00563, 00566, 00567, 00580, 00600, 00604, 00620, 00625, 00626, 00630, 00632, 00635, 00640, 00670, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00754, 00756, 00770, 00790, 00792, 00794, 00796, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00840, 00842, 00844, 00846, 00848, 00851, 00860, 00862, 00864, 00865, 00866, 00868, 00870, 00872, 00873, 00880, 00882, 00902, 00904, 00906, 00908, 00910, 00912, 00914, 00916, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00932, 00934, 00936, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01130, 01140, 01150, 01160, 01170, 01173, 01200, 01202, 01210, 01212, 01214, 01215, 01220, 01230, 01232, 01234, 01250, 01260, 01270, 01272, 01274, 01320, 01340, 01360, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01420, 01430, 01432, 01440, 01442, 01444, 01462, 01464, 01470, 01472, 01474, 01480, 01482, 01484, 01486, 01490, 01500, 01502, 01520, 01522, 01610, 01620, 01622, 01630, 01634, 01636, 01638, 01650, 01652, 01654, 01656, 01670, 01680, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01756, 01758, 01760, 01770, 01772, 01780, 01782, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01916, 01920,

2026 AQR QCDR Measure Specifications Manual

01922, 01924, 01925, 01926, 01930, 01931, 01932, 01933, 01937, 01938, 01939, 01940, 01941, 01942, 01951, 01952, 01958, 01960, 01961, 01962, 01963, 01965, 01966, 01967, 01991, 01992

AND

Place of service code 19, 21, 22, or 24

AND

Patient is taking methadone as an outpatient or has a prescription for methadone in their admittance documentation. **12A27**

Denominator Exclusions:

None

NUMERATOR (Measure B)

All patients where methadone is either continued or changed to an equivalent IV dose to maintain the therapeutic level regimen during the perioperative period or, if in an ambulatory setting, was provided as scheduled prior to surgery.

***Numerator Note:** In the inpatient setting, Methadone may either be continued or changed to an equivalent IV dose to maintain therapeutic levels. Documentation of methadone medication provided as scheduled prior to surgery in the ambulatory setting is acceptable for this measure.*

***Numerator Note:** For this measure, the perioperative period includes preoperative, intraoperative, and postoperative period. The postoperative period ends, for this measure, when the patient is discharged from the post-anesthesia care unit or recovery location.*

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

Methadone therapy for the patient is continued or changed to an equivalent IV dose to maintain the therapeutic level regimen during the perioperative period or, if in an ambulatory setting, methadone was provided as scheduled prior to surgery.

OR

Denominator Exception:

Documentation that patient received instruction from their surgical, anesthesia, and addiction treatment provider to stop methadone prior to inpatient or ambulatory procedure. Documentation of medical reasons for not continuing or changing to an equivalent IV dose of methadone treatment during the perioperative period or, if in an ambulatory setting, methadone was not provided as scheduled prior to surgery. Medical reasons may include patient transfer to intensive care unit, a patient diagnosis of hypotension or shock, unstable plasma levels, or patient is unable to tolerate the daily methadone regimen.

OR

Performance Not Met:

Methadone therapy for the patient is NOT continued or changed to an equivalent IV dose to maintain the therapeutic level regimen during the perioperative period or, if in an ambulatory setting, methadone was NOT provided as scheduled prior to surgery.

CBE Number: Not applicable

eCQM: Not applicable

Data Source: Administrative claims data; Claims; EHR; Paper medical record; Record review; Registry

Care Setting: Ambulatory; Ambulatory Surgical Center; Hospital; Hospital Inpatient; Hospital Outpatient; Office Based Surgery Center

2026 AQR QCDR Measure Specifications Manual

Telehealth: No

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number of Performance Rates: 2 – Average

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Ratio Measure: No

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI81: Use of Peripheral Nerve Blocks for Pain Management in Hip Fracture Patients

Provision licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description:

All patients, aged 65 and older, who are diagnosed with a hip fracture and receive peripheral nerve blocks for pain management in the emergency room or inpatient setting.

NQS Domain/Meaningful Measures Area

Patient Safety

Measure Type

Process

High Priority Status

Yes

Inverse Measure

No

Instructions:

This measure is to be reported when a patient aged 65 and older, diagnosed with an isolated hip fracture, receives peripheral nerve blocks for pain management in the emergency and/or inpatient settings of the hospital. It is anticipated that qualified anesthesia providers and eligible clinicians who provide care for the denominator-eligible population will submit this measure.

Measure Submission Type:

Measure data may be submitted by individual MIPS eligible clinicians, groups, or third-party intermediaries via the Qualified Clinical Data Registry option.

Measure Reporting via the Qualified Clinical Data Registry:

CPT (Current Procedural Terminology) codes, and patient demographics and registry codes are used to identify patients who are included in the measure denominator.

Denominator

Patients, aged 65 and older, with a diagnosis of an isolated hip fracture.

Denominator Criteria (Eligible Cases):

All patients aged 65 and older

AND

Diagnosis of isolated hip fracture: 12A41

AND

Patient encounter during the reporting period (CPT):

64450, 64447, 64448, 64449, 62323, 64474

AND

Place of service code: 21, 23

2026 AQR QCDR Measure Specifications Manual

Denominator Exclusions

Emergent Cases: M1142

Multiple surgical procedures: 12A42

Numerator

Patient received peripheral nerve blocks for pain management in the emergency room and/or inpatient setting.

Numerator Note: For the purposes of this measure, the numerator is met if a peripheral nerve block via a single injection and/or catheter is administered for pain management during the hospital care episode.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

12A43 Peripheral nerve block was used for pain management

OR

Denominator Exception:

12A44 Documentation of patient reason(s), process reason(s) or medical reason(s) for not receiving peripheral nerve block (i.e., medical contraindication, patient has allergy to local anesthetics, patient has coagulopathy or ongoing anticoagulation without appropriate reversal, infection at the PNB insertion site, patient refusal), or an anesthesiologist was not consulted.

OR

Performance Not Met:

12A45 Peripheral nerve block was NOT used for pain management

CBE Number: Not applicable

eCQM: Not applicable

Data Source: Claims/Paper Medical Record, Registry

Care Setting: Hospital Inpatient

Telehealth: No

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number of Performance Rates: 1

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Ratio Measure: No

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS

2026 AQR QCDR Measure Specifications Manual

Measure Title

AQI82: Intraoperative Antibiotic Redosing in Open Cardiac Surgeries

Provation licensed this measure from Anesthesia Quality Institute (AQI) National Anesthesia Clinical Outcomes Registry (NACOR) in Collaboration with American Society of Anesthesiologists (ASA)

Measure Description:

Percentage of patients who undergo an open cardiac procedure over 240 minutes duration and who receive initial cephalosporin or clindamycin administration within 60 minutes before surgical incision and receive either a bolus infusion or an intravenous cephalosporin antibiotic re-dose at four-hour intervals.

NQS Domain/Meaningful Measures Area

Patient Safety

Measure Type

Process

High Priority Status

Yes

Inverse Measure

No

Instructions:

This measure is to be reported each time a patient undergoes an open cardiac surgical procedure for over 240 minutes (4 hours) during the reporting period. The measure time commences 60 to 120 minutes before surgical incision during which the patient receives their initial antibiotic dose. It is anticipated that qualified anesthesia professionals and eligible clinicians who provide denominator-eligible services will submit this measure.

Measure Submission Type:

Measure data may be submitted by individual MIPS-eligible clinicians, groups, or third-party intermediaries via the Qualified Clinical Data Registry option.

Measure Reporting via the Qualified Clinical Data Registry:

CPT (Current Procedural Terminology) codes, denominator criteria, registry codes, and patient demographics are used to identify patients who are included in the measure denominator. The numerator options included in this specification are used to submit the quality actions as allowed by the measure.

Denominator

All patients who undergo an open cardiac procedure over 240 minutes (4 hours) duration.

Denominator Criteria (Eligible Cases):

All patients

AND

Procedure greater than 240 minutes (4 hours) duration: **12A02**

AND

Patient encounter during the reporting period (CPT):

00560, 00561, 00562, 00563, 00566, 00567, 00580

Denominator Exclusions

Non-cardiac, transcatheter/endovascular, electrophysiology/catheterization, and other cardiac cases (determined by the procedure type). **12A03**

2026 AQR QCDR Measure Specifications Manual

- Non-cardiac: 44950, 44955, 44960, 47562, 47563, 47564, 27447, 61510, 42820, 42821, 42825, 42826, 55866, 58150, 58260–58294, 8541–85573.
- Transcatheter/Endovascular: 37220, 37221, 37224, 37226, 34802, 37184, 37187.
- Electrophysiology/Catheterization: 93619, 93620, 93621, 93622, 93653, 93654, 93656, 93657, 93609.
- Other Cardiac Cases: 35301, 35302, 365555, 36557, 365558, 36560, 36555, 36561, 36568, 36569, 36572, 36573, 36556, 36555, 36557, 36558, 36560, 36561, 36568, 36569, 36572, 36620, 37221, 3724, 37226, 37228, 92920, 92928, 92928, 34001, 34101, 34111, 34201, 34203, 37184, 37185, 37186, 37187.
- Chronic kidney disease. N18.1, N18.2, N18.3, N18.30, N18.31, N.18.32, N18.4, N 18.5, N18.6, N18.22, E11.22, I12.x, I13.x, Z99.2.

Numerator

Patients who receive administration of initial cephalosporin antibiotic prophylaxis within 60 minutes before surgical incision and either bolus infusion or intravenous redosing of their cephalosporin within four hours of initial cephalosporin antibiotic prophylaxis.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

12A46 Patients received administration of initial cephalosporin antibiotic prophylaxis within 60 minutes before surgical incision administration and received either bolus infusion or intravenous redosing cephalosporin within 240 minutes (4 hours) of initial cephalosporin antibiotic prophylaxis or bolus infusion.

OR

Denominator Exception:

12A47 Documentation of patient reason(s) for not receiving antibiotic redosing (i.e., medical contraindication; patient already received scheduled antibiotics).

OR

Performance Not Met:

12A48 Patients did NOT receive administration of initial cephalosporin antibiotic prophylaxis within 60 minutes before surgical incision and/or did not receive administration of a bolus infusion or intravenous redosing of their cephalosporin within 240 minutes (4 hours) of initial cephalosporin antibiotic prophylaxis.

CBE Number: Not applicable

eCQM: Not applicable

Data Source: Claims/Paper Medical Record, Registry

Care Setting: Hospital Inpatient

Telehealth: No

Measure Steward: American Society of Anesthesiologists (ASA)/Anesthesia Quality Institute (AQI)

Number of Performance Rates: 1

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Ratio Measure: No

Risk Adjustment: No

2026 AQR QCDR Measure Specifications Manual

MIPS Reporting Option: Traditional MIPS

2026 AQR QCDR Measure Specifications Manual

Measure Title

ABG44: Low Flow Inhalational General Anesthesia

Provision licensed this measure from ABG QCDR

Measure Description:

Percentage of patients aged 18 years or older, who undergo an elective procedure lasting 30 minutes or longer requiring inhalational general anesthesia who during the maintenance phase of the anesthetic have a total fresh gas flow less than or equal to 1 L/min (less than or equal to 2 L/min for Sevoflurane).

Measure Type

Efficiency

High Priority Status

Yes

Inverse Measure

No

Instructions

This measure is to be reported each time a patient undergoes an elective procedure in which inhalational general anesthesia is used. It is anticipated that qualified anesthesia providers and eligible clinicians who provide denominator-eligible services will submit this measure.

Measure Reporting via the Qualified Clinical Data Registry

Patient demographics and CPT codes are used to identify patients who are included in the measure denominator. G-codes and Registry Codes are used to capture the numerator.

Denominator

All patients aged 18 years or older, who undergo an elective procedure lasting 30 minutes or longer requiring inhalational general anesthesia. **(1095)**

Denominator Criteria (Eligible Cases):

Patients aged 18 years and older

AND

Elective procedure

AND

Patient who receives inhalational general anesthesia

AND

Procedure lasts 30 minutes or longer

Patient encounter during the reporting period (CPT) 00100, 00102, 00103, 00104, 00120, 00124, 00126, 00140, 00142, 00144, 00145, 00147, 00148, 00160, 00162, 00164, 00170, 00172, 00174, 00176, 00190, 00192, 00210, 00211, 00212, 00214, 00215, 00216, 00218, 00220, 00222, 00300, 00320, 00322, 00350, 00352, 00400, 00402, 00404, 00406, 00410, 00450, 00454, 00470, 00472, 00474, 00500, 00520, 00522, 00524, 00528, 00529, 00530, 00532, 00534, 00537, 00539, 00540, 00541, 00542, 00546, 00548, 00550, 00560, 00566, 00580, 00600, 00604, 00620, 00625, 00626, 00630, 00632, 00635, 00640, 00670, 00700, 00702, 00730, 00731, 00732, 00750, 00752, 00754, 00756, 00770, 00790, 00792, 00794, 00796, 00797, 00800, 00802, 00811, 00812, 00813, 00820, 00830, 00832, 00840, 00842, 00844, 00846, 00848, 00851, 00860, 00862, 00864, 00865, 00866, 00868, 00870, 00872, 00873, 00880, 00882, 00902, 00904, 00906, 00908, 00910, 00912, 00914, 00916, 00918, 00920, 00921, 00922, 00924, 00926, 00928, 00930, 00932, 00934, 00936, 00938, 00940, 00942, 00944, 00948, 00950, 00952, 01112, 01120, 01130, 01140, 01150, 01160, 01170, 01173, 01200, 01202,

2026 AQR QCDR Measure Specifications Manual

01210, 01212, 01214, 01215, 01220, 01230, 01232, 01234, 01250, 01260, 01270, 01272, 01274, 01320, 01340, 01360, 01380, 01382, 01390, 01392, 01400, 01402, 01404, 01420, 01430, 01432, 01440, 01442, 01444, 01462, 01464, 01470, 01472, 01474, 01480, 01482, 01484, 01486, 01490, 01500, 01502, 01520, 01522, 01610, 01620, 01622, 01630, 01634, 01636, 01638, 01650, 01652, 01654, 01656, 01670, 01680, 01710, 01712, 01714, 01716, 01730, 01732, 01740, 01742, 01744, 01756, 01758, 01760, 01770, 01772, 01780, 01782, 01810, 01820, 01829, 01830, 01832, 01840, 01842, 01844, 01850, 01852, 01860, 01924, 01925, 01926, 01930, 01931, 01932, 01933, 01935, 01936, 01951, 01952, 01961, 01962, 01963, 01965, 01966

Denominator Exceptions

Patient or technical reason exists for not providing low flow inhalational anesthesia (e.g., flow meter not capable of generating low flows, patient hypermetabolic, lack of CO₂ absorbents without KOH and low concentrations of NaOH, etc.) **(1096)**

Numerator

Patients who undergo an elective procedure lasting 30 minutes or longer requiring inhalational general anesthesia who during the maintenance phase of the anesthetic have a total fresh gas flow less than or equal to 1 L/min (less than or equal to 2 L/min for Sevoflurane).

Numerator Definition

Inhalational general anesthesia is defined as the use of at least one inhalational anesthetic gas (e.g., halothane, isoflurane, desflurane, sevoflurane, nitrous oxide) as the primary mode of anesthesia for the procedure.

The maintenance phase of the inhalational anesthetic is defined as the portion of the case in which Stage III surgical anesthesia (e.g., unconsciousness, amnesia, immobility, unresponsive to surgical stimulation) is achieved at a safe anesthetic depth while also maintaining respiratory and hemodynamic stability. This occurs between the induction and emergence phases of the case.

Fresh gas flow (FGF) is defined as the combined admixture of medical gases such as air, oxygen, or nitrous oxide as well as volatile anesthetics as set by the anesthesia provider.

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

The total FGF is reduced to less than or equal to 1 L/min (less than or equal to 2 L/min for Sevoflurane) for the duration of the maintenance phase of the anesthetic **(1097)**.

OR

Performance Not Met:

The total FGF is greater than 1 L/min (greater than 2 L/min for Sevoflurane) for the duration of the maintenance phase of the anesthetic **(1098)**.

CBE Number: Not Applicable

eCQM: Not Applicable

Data Source: Hybrid; Other: Other: Medical Record, Registry

Care Setting: Ambulatory Surgical Center; Imaging Facility; Office Based Surgery Center; Ambulatory Care: Hospital; Hospital Inpatient; Hospital Outpatient

Telehealth: No

Measure Steward: ABG QCDR in Collaboration with Insight Medical Data Services, LLC

2026 AQR QCDR Measure Specifications Manual

Number of Multiple Performance Rates: Not Applicable

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS, MVP

2026 AQR QCDR Measure Specifications Manual

Measure Title

ABG45: Aspiration prevention in patients with gastric distension

Provision licensed this measure from ABG QCDR

Measure Description:

Percentage of patients **18 years** and older with a current diagnosis of gastrointestinal obstruction, ileus, incarcerated hernia, or patients taking GLP-1 receptor agonists, or patients with gastroparesis, who undergo a surgical procedure under anesthesia and are treated preoperatively with a mitigation strategy that reduces the risk of aspiration during the surgical procedure.

Measure Type

Process

High Priority Status

Yes

Inverse Measure

No

Instructions

This measure is to be reported each time an adult patient with one or more of the following conditions undergoes a surgical procedure under anesthesia:

- a) has gastrointestinal obstruction, ileus, or an incarcerated hernia
- b) is taking a GLP-1 receptor agonists
- c) has gastroparesis

Denominator

All patients aged 18 years or older who undergo a surgical procedure under anesthesia and have one or more of the following conditions:

- a) has gastrointestinal obstruction, ileus, or an incarcerated hernia
- b) is taking a GLP-1 receptor agonists
- c) has gastroparesis

Denominator Criteria (Eligible Cases)

Patients aged 18 years and older

AND

Current diagnosis of gastrointestinal obstruction, ileus or incarcerated hernia

OR

Has gastroparesis

OR

Is taking GLP-1 agonists (last dose < 4 half-lives)

AND

Receives anesthesia (MAC, regional, general, neuraxial)

Patient encounter during the reporting period (CPT): All cases

Denominator Exclusion

2026 AQR QCDR Measure Specifications Manual

- Local anesthesia only (no sedation)
- Patient already intubated

Numerator

Patients with current diagnosis of gastrointestinal obstruction, ileus or incarcerated hernia or are patients taking GLP-1 receptor agonists or are patients with gastroparesis, who prior to the surgical procedure under anesthesia have one of the following mitigation strategies applied:

1. Clinical imaging demonstrating empty stomach contents (≤ 1.5 ml/kg of fluid and no solids) within one hour of procedure (MRI/CT/gastric ultrasound)
2. Evacuation of stomach contents by naso/orogastric tube within one hour of procedure
3. Awake intubation
4. Maintenance of airway reflexes throughout the case

Numerator Definition

Numerator Quality-Data Coding Options for Reporting Satisfactorily

Performance Met:

45PM one or more of the four mitigation strategies applied prior to the procedure

OR

Denominator Exception:

45DE Patients taking GLP-1 RAs or have gastroparesis but do not display symptoms of gastric distension

- Symptomatic gastric distension may be defined as patients who exhibits two of the three following symptoms:
 - 1) nausea and/or vomiting
 - 2) dyspepsia/abdominal pain
 - 3) abdominal bloating/distension

OR

Performance Not Met:

45PNM none of the four mitigation strategies applied prior to the procedure

CBE Number: Not Applicable

eCQM: Not Applicable

Data Source: Administrative claims data; EHR

Care Setting: All Settings

Telehealth: No

Measure Steward: ABG QCDR

Number of Multiple Performance Rates: Not Applicable

Proportion Measure Scoring: Yes

Continuous Measure Scoring: No

Ratio Measure: No

2026 AQR QCDR Measure Specifications Manual

Risk Adjustment: No

MIPS Reporting Option: Traditional MIPS